

# Leukaemia Section

## Short Communication

### del(13q) in ALL

David Betts

Department of Oncology, University Children's Hospital, Steinwiesstr. 75, CH-8032 Zürich, Switzerland (DB)

Published in Atlas Database: November 2004

Online updated version : <http://AtlasGeneticsOncology.org/Anomalies/del13qALLID1188.html>

DOI: 10.4267/2042/38154

This work is licensed under a Creative Commons Attribution-Noncommercial-No Derivative Works 2.0 France Licence.

© 2005 Atlas of Genetics and Cytogenetics in Oncology and Haematology

## Identity

### Note

Deletions of chromosome 13q are a non-random finding in a broad spectrum of haematological neoplasms, including B-cell chronic lymphocytic leukemia (CLL), non-Hodgkin's lymphoma (NHL) and multiple myeloma (MM) and acute myeloid leukaemia (AML).

## Clinics and pathology

### Disease

Acute lymphoblastic leukaemia (ALL).

### Phenotype/cell stem origin

No specific immunophenotype observed.

### Epidemiology

A del(13q) chromosome is found in approximately 2% of cases in both adult and childhood disease at presentation. Up to 4% of cases may have some loss of 13q material, either through full monosomy or unbalanced rearrangements. Incidence of chromosome 13 deletions is higher at relapse.

### Prognosis

May confer an increased risk of treatment failure but to date has not been shown to be an independent prognostic indicator.

## Cytogenetics

### Cytogenetics morphological

Various breakpoints reported. The centromeric breakpoint is typically in the 13q12-14 region and telomeric between 13q21 and 13qter. Loss of all or part of 13q14 is common to almost all cases. Occurs as a

sole event in approximately 10% of cases. There are also rare reports of translocations also leading to a partial 13q deletion. Monosomy 13 is also reported but occurs very rarely as a sole aberration. Under representation of chromosome 13 is often found in hypotriploid cases.

### Additional anomalies

Most cases with del(13q) will have additional aberrations, but there is no consistent picture and the events can include the typical non-random events in ALL.

## Genes involved and proteins

### Note

Critical region in 13q14 appears to lie telomeric to RB1.

## References

Chung CY, Kantarjian H, Haidar M, Starostik P, Manshouri T, Gidel C, Freireich E, Keating M, Albitar M. Deletions in the 13q14 locus in adult lymphoblastic leukemia: rate of incidence and relevance. *Cancer*. 2000 Mar 15;88(6):1359-64

Heerema NA, Sather HN, Sensel MG, Lee MK, Hutchinson RJ, Nachman JB, Reaman GH, Lange BJ, Steinherz PG, Bostrom BC, Gaynon PS, Uckun FM. Abnormalities of chromosome bands 13q12 to 13q14 in childhood acute lymphoblastic leukemia. *J Clin Oncol*. 2000 Nov 15;18(22):3837-44

Schneider NR, Carroll AJ, Shuster JJ, Pullen DJ, Link MP, Borowitz MJ, Camitta BM, Katz JA, Amylon MD. New recurring cytogenetic abnormalities and association of blast cell karyotypes with prognosis in childhood T-cell acute lymphoblastic leukemia: a pediatric oncology group report of 343 cases. *Blood*. 2000 Oct 1;96(7):2543-9

Cavé H, Avet-Loiseau H, Devaux I, Rondeau G, Boutard P, Lebrun E, Méchinaud F, Vilmer E, Grandchamp B. Deletion of chromosomal region 13q14.3 in childhood acute lymphoblastic leukemia. *Leukemia*. 2001 Mar;15(3):371-6

Kovacs BZ, Niggli FK, Betts DR. Aberrations involving 13q12 approximately q14 are frequent secondary events in childhood acute lymphoblastic leukemia. *Cancer Genet Cytogenet.* 2004 Jun;151(2):157-61

---

*This article should be referenced as such:*

Betts D. *del(13q) in ALL.* *Atlas Genet Cytogenet Oncol Haematol.* 2005; 9(1):24-25.

---